



SCIENCE
AND DENTISTRY.
BONDED.



PRODUCT SPOTLIGHT



Table of Contents

- **Educational Resources**
 - eBooks Page 3
- **Adhesives**
 - All-Bond Universal® Page 4
- **Cements**
 - Duo-Link Universal™ Page 5
- **Etchants**
 - Select HV® Etch w/BAC, Uni-Etch® w/BAC, & Etch-37™ w/BAC..... Page 6
- **TheraFamily**
 - TheraCal LC® Page 7
 - TheraCal PT® Page 8
 - TheraCem® Page 9
 - TheraBase® Page 10
- **Primers**
 - Z-Prime™ Plus..... Page 11
- **Supportive Products**
 - ZirClean® Page 12
- **Core Materials**
 - Core-Flo™ DC, & Core-Flo™ DC Lite Page 13
- **Varnish**
 - FluoroCal™ Page 15
- **Techniques**
 - Self-Etch, Total-Etch, and Selective-Etch Bonding Page 16
 - Pulp Exposures (Direct Pulp Capping) Page 17
 - Deep Preparations (Indirect Pulp Capping) Page 17
 - Sandwich Technique Page 18
 - Pulpotomy..... Page 18
 - Core Build-ups Page 19
 - Intraoral Repairs of Porcelain (PFM or PFZr) Page 20
 - Conventional Cementation of Crowns (Retentive Preps)..... Page 21
 - Bonded Cementation of Inlays/Onlays, Bridges, Crowns (Short/Tapered Preps)..... Page 22
 - Veneer Cementation..... Page 23
- **Quick Guides**
 - Zirconia Bonding & Sandwich Technique..... Page 24
 - Veneer Cementation & Cementation With a Bonding Agent Page 25
 - Cementation Without a Bonding Agent & Post Cementation With Core Build-up Page 26

Visit us at www.bisco.com for a complete product catalog!



At BISCO “adhesion is our passion,” and we dedicate ourselves to understanding and improving the ability to bond restorations. Our goal is to make your life easier, while helping you perform your best dentistry! We place tremendous value on sharing our research and scientific knowledge to benefit you and your practice, and offering award-winning products that provide reliable solutions to your clinical needs.

With BISCO, you are no longer alone. If you have a question about restorative protocols or are wondering which products to use (or maybe you are just looking for someone to talk to), give us a call. We’re here to help!




BISCO HEADQUARTERS
SCHAUMBURG, ILLINOIS



DR. BYOUNG SUH
FOUNDER AND PRESIDENT OF BISCO
- CHEMIST
- AUTHOR OF PRINCIPLES OF
ADHESION DENTISTRY
- EXCELLENT GOLFER


Restorative Solutions eBook

Meet the Award-Winning TheraFamily




Understanding Cements eBook

A Guide From Prep to Post-Op




Universal Adhesives eBook

A Brief Clinical Overview



The ABCs of Zirconia Bonding

Bonding to Zirconia is Achievable



*USE YOUR MOBILE DEVICE
TO SCAN THE QR CODES
AND LEARN MORE!*

All-Bond Universal®

Light-Cured Dental Adhesive

All-Bond Universal is the culmination of over 30 years of adhesive research at BISCO. All-Bond Universal is compatible with all light-, self-, and dual-cured resin composite and cement materials for all direct and indirect procedures.



Versatile
Offers the flexibility for total-, self-, or selective-etch procedures



MDP
Contains MDP for enhanced durability



High Shear Bond Strengths
High shear bond strength to all substrates



Low Film Thickness
Less than 10 microns



Single Bottle System
Other adhesives may need more than one bottle for indirect restorations, but with All-Bond Universal, NO activator is required



Total "Universality"
Compatible with all light-, self- and dual-cured resin composite and cement materials for all direct and indirect procedures

Dentistry courtesy of Tyler Lasseigne, DDS, CDT



1. Apply Z-Prime Plus



2. Apply All-Bond Universal



3. Apply Duo-Link Universal

LEARN MORE ABOUT ALL-BOND UNIVERSAL



Order Info: Bottle

All-Bond Universal Standard Kit	B-72020K
All-Bond Universal (6ml)	B-7202P

Unit-Dose Packages

All-Bond Universal Unit-Dose (50pk)	B-73050K
All-Bond Universal Unit-Dose (100pk)	B-73100K

*Universality: The powerful feeling of the entire universe in one single bottle.





Duo-Link Universal™ Resin Luting Cement

Duo-Link Universal is specially formulated for cementation of ALL* indirect restorations. The adhesive resin cement is intended for use with adhesives designed for compatibility with all dental materials, including all BISCO adhesives.



Easy Clean-Up
Formulated to allow for quick and easy removal of excess cement



Easy to Use
Auto-mix, dual-syringe provides a consistent mix and easy placement



Radiopaque
Visible on radiograph to easily distinguish from caries



CAD/CAM Restorations
Ideal for all chairside and lab-fabricated restorations




High Degree of Conversion
In both light- and self-cured modes ensures a strong, long lasting restoration



Universal For All Cementation Procedures*
Crowns, bridges (fixed prosthesis), inlays, onlays, and posts/dowels, fabricated from metal, composite, porcelain, ceramic, zirconia, alumina, CAD/CAM restorations, etc.

Dentistry courtesy of Darren D. Simpson, DDS



1. Duo-Link Universal dispensed in the crown



2. All-Bond Universal applied, air dried, and light-cured



3. Restoration seated. Excess cement removed



4. Final restoration

LEARN MORE ABOUT
DUO-LINK UNIVERSAL



Order Info: System Kit with All-Bond Universal B-19620K

Universal Dual-Syringe (8g) A-19030P
Milky White Dual-Syringe (8g) A-197MWP

* It is recommended to use BISCO's CHOICE™ 2 for veneer cementation.





Select HV[®] Etch w/BAC Uni-Etch[®] w/BAC Etch-37[™] w/BAC

Phosphoric Acid Etchants with Benzalkonium Chloride (BAC)

Select HV Etch is a 35% high-viscosity phosphoric acid etchant with BAC. It is specifically formulated for pin-point placement to etch enamel when using an adhesive in selective-etch mode.

Uni-Etch w/BAC and Etch-37 w/BAC are 32% and 37% semi-gel phosphoric acid etchants with BAC. They are specifically formulated to be easily applied to larger surface areas and rinsed cleanly with no residue.



Blue Color
Easy visualization and contrast



Contains BAC
In-vitro research shows benzalkonium chloride is effective against Streptococcus mutans^{1,2}

NOTE: Inclusion of BAC has not been shown to correlate with a reduction in secondary decay in patients. In vivo clinical studies to evaluate the effects of BAC on oral bacteria or caries have not been performed.



Easy Clean Up
Rinses away cleanly and quickly leaving no residue to interfere with bonding

Dentistry courtesy of Michael Morgan, DDS



Select HV Etch w/BAC demonstrating pin-point placement on enamel

LEARN MORE ABOUT
ETCHANTS



Order Info: Select HV Etch w/BAC (35%)

- 30ml Bulk Syringe Kit E-59200K
- Bulk Syringe Refill (30ml) E-59160P
- 4 Syringe Package (5g ea.) E-59110P

Etch-37 w/BAC (37%)

- Bulk Bottle (30g) E-5638EB
- Bulk Syringe Refill (30ml) E-56741P
- 4 Syringe Package (5g ea.) E-5503EBM

Uni-Etch w/BAC (32%)

- Bulk Bottle (30g) E-5637EB
- Bulk Syringe Refill (30ml) E-56621P
- 4 Syringe Package (5g ea.) E-5502EBM

1. M.Sc.Dt. Emre ÖZEL, Dr. Haktan YURDAGÜVEN, Yrd.Doç.Dr. Esra CAN SAY, Prof.Dr. Sesin KOCAGÖZ, Evaluation of the Antibacterial Activity of Disinfectant Solutions with Phosphoric Acids Against Streptococcus Mutans. Journal of Hacettepe Faculty of Dentistry, Volume: 29, Issue 4, Page: 8-14, 2005
2. M. TURKUN1, Z. ERGUCU, L.S. TURKUN, E.U. CELIK, and M. ATEŞ, Is Phosphoric Acid Sufficiently Antibacterial? J Dent Res 85 (Spec Iss B):abstract number 1605, 2006 (www.dental.research.org).



TheraCal LC® Resin-Modified Calcium Silicate Pulp Protectant/Liner

TheraCal LC is a light-cured resin modified calcium silicate ideal for direct and indirect pulp capping and as a protective liner.



Calcium Release*
Stimulates hydroxyapatite formation^{1,2} and secondary dentin bridge formation^{2,3}



Radiopaque
Visible on radiograph to easily distinguish from caries



Moisture Tolerant
Has low solubility and will not wash out over time



Alkaline pH
Alkaline pH promotes pulp vitality³



Insulates Pulp
Forms a protective barrier that insulates the pulp^{4,5}

Dentistry courtesy of Ross Nash, DDS



1. Distal-occlusal caries present on an asymptomatic mandibular first premolar



2. Incomplete excavation of caries leaving affected dentin without exposing the pulp



3. Placement of TheraCal LC on moist dentin



4. Place 1mm of TheraCal LC and light cure for 20 seconds

LEARN MORE ABOUT
THERACAL LC



OR VISIT
THERAFAMILY.COM

Order Info: TheraCal LC 4-Syringe Pack
TheraCal LC Syringe (1g)

H-33014P
H-3301P

* BISCO has, on file, the calcium release data for TheraCal LC

1. Gandolfi MG, Siboni F, Prati C. Chemical-physical properties of TheraCal, a novel light-curable MTA-like material for pulp capping. International Endodontic Journal. 2012 Jun;45(6):571-9.
2. ADA definitions for direct and indirect pulp capping at: www.ada.org/en/publications/cdt/glossary-of-dental-clinical-and-administrative-ter
3. T. Okabe, M. Sakamoto, H. Takeuchi, K. Matsushima. Effects of pH on Mineralization Ability of Human Dental Pulp Cells. Journal of Endodontics. Volume 32, Number 3, March 2006.
4. Sangwan P; Sangwan A; Duhan J; Rohilla A. Tertiary dentinogenesis with calcium hydroxide: a review of proposed mechanisms. Int Endod J. 2013; 46(1):3-19
5. Selcuk SAVAS, Murat S. BOTSALI, Ebru KUCUKYILMAZ, Tugrul SARI. Evaluation of temperature changes in the pulp chamber during polymerization of light-cured pulp-capping materials by using a VALO LED light curing unit at different curing distances. Dent Mater J. 2014;33(6):764-9.





TheraCal PT®

Dual-Cured Resin-Modified Calcium Silicate Pulpotomy Treatment

TheraCal PT is a biocompatible, dual-cured, resin-modified calcium silicate designed for pulpotomy treatment. TheraCal PT maintains tooth vitality by performing as a barrier and protectant of the dental pulpal complex.



Calcium Release
Unique hydrophilic matrix facilitates calcium release



Radiopaque
Visible on radiograph to easily distinguish from caries



Easy Syringe Application
Manual mixing is not required, the dispensing tip creates a uniform mix allowing for direct placement



Alkaline pH
Alkaline pH promotes pulp vitality¹



Dual Cured
One-layer confidence



Moisture Tolerant
Low water solubility

Dentistry courtesy of Juan Carlos Hernández Cabanillas, DDS



1. Perform pulpotomy treatment and achieve hemostasis



2. Place TheraCal PT directly in the pulp chamber, and ensure good adaptation to the cavity walls and margins



3. After light curing TheraCal PT for 10 seconds, place desired adhesive, base, and/or restoration following manufacturer's directions

LEARN MORE ABOUT
THERACAL PT



OR VISIT
THERAFAMILY.COM

Order Info: TheraCal PT Dual-Syringe (4g) H-34110P

1. T. Okabe, M. Sakamoto, H. Takeuchi, K. Matsushima. Effects of pH on Mineralization Ability of Human Dental Pulp Cells. Journal of Endodontics. Volume 32, Number 3, March 2006.





TheraCem[®] Self-Adhesive Resin Cement

TheraCem is a dual-cured, calcium- and fluoride-releasing, self-adhesive resin cement indicated for luting crowns, bridges, inlays, onlays and posts (prefabricated metal/non-metal/fiber posts).



Calcium and Fluoride Release

TheraCem offers continuous calcium and fluoride release*



Radiopaque

Visible on radiograph to easily distinguish from caries



Alkaline pH

TheraCem transitions from acidic to alkaline pH in minutes*



Easy Clean-Up

Specially formulated to allow for quick and easy clean-up



High Degree of Conversion

A high degree of conversion ensures a higher physical strength



MDP

TheraCem contains MDP offering a strong bond to zirconia, metal, and alumina substrates with no priming or etching required

Dentistry courtesy of Joseph Kim DDS, JD, FAGD, FICOI



1. Prepare tooth.



2. Seat restoration with TheraCem.



3. Remove excess cement.



4. Inspect margins.

LEARN MORE ABOUT
THERACEM



OR VISIT
THERAFAMILY.COM

Order Info: Natural Dual-Syringe (8g)

D-46311P

* Data on file. BISCO, Inc.





TheraBase®

Self-Adhesive Calcium Releasing Base/Liner

TheraBase is a dual-cure, calcium and fluoride-releasing, self-adhesive base/liner. It is the ideal dual-cured material that will polymerize even in deep restorations where light cannot reach.



Calcium and Fluoride Release

TheraBase offers continuous calcium and fluoride release*



High Flexual Strength

Stronger and more fracture resistant



High Compressive Strength

Absorbs shock and stress from occlusal forces without fracturing



Radiopaque

Visible on radiograph to easily distinguish from caries



Alkaline pH

Generates an alkaline pH in minutes, which promotes pulp vitality¹



High Degree of Conversion

Ensures enhanced physical properties

Dentistry courtesy of Dr. Rual Euan, DDS



1. After cavity preparation, all water was removed using a stream of air, leaving the surface visibly moist. TheraCal LC was applied on the small pulp exposure and light-cured for 20 seconds.



2. TheraBase was applied to the dentin surface of the prepared cavity directly from the dispensing syringe.



3. TheraBase was light-cured for 20 seconds. If desired, TheraBase can be allowed to self-cure for 4 minutes.



4. A selective-etch bonding technique was used to condition the surface of the preparation. Any bonding technique can be applied.



5. All-Bond Universal was applied following manufacturer's instructions.



6. Restorations were filled with a light-cure composite material following manufacturer's instructions.

LEARN MORE ABOUT
THERABASE



OR VISIT
THERAFAMILY.COM

Order Info: TheraBase Single Syringe Pack H-35001P

* Data on file. BISCO, Inc.

1. T. Okabe, M. Sakamoto, H. Takeuchi, K. Matsushima. Effects of pH on Mineralization Ability of Human Dental Pulp Cells. Journal of Endodontics. Volume 32, Number 3, March 2006.



Z-Prime™ Plus

Zirconia - Alumina - Metal Primer

Z-Prime Plus is a one-bottle primer used to enhance adhesion between indirect restorative materials and composite resin cements.



High Bond Strength

To zirconia, alumina, and metal restorations



Versatile

Can be used with all metal and metal-oxide substrates



Compatibility

Compatible with light-cured and dual-cured resin luting cements



Enhances Bond Strength

Significantly enhances bond strengths of other resin cements*



Single Bottle

Convenient single bottle delivery offers ease of dispensing



MDP

Contains MDP, allowing for a strong bond to zirconia, metal, and alumina substrates

Order Info: 1 Bottle Z-Prime Plus (2ml)
1 Bottle Z-Prime Plus (4ml)

B-6002P
B-6001P

* Data on file. BISCO, Inc.

LEARN MORE ABOUT
Z-PRIME PLUS





ZirClean® Restoration Cleaner

ZirClean is an extraoral cleaning agent designed for the non-abrasive cleaning of the bonding surfaces of zirconia, ceramic and metal restorations, after intraoral try-in.



Easy to Use

Syringe delivery system offers improved ease of use



Easy Clean up

Easy clean-up of the material



Contaminant Remover

Contaminants such as saliva can be removed to help achieve better bond strengths after try-in*



Reliable Results

Helps to achieve reliable cementation results with zirconia restorations after intraoral try-in*

Order Info: 1 Syringe ZirClean (5g)

B-7351P

* As compared to untreated samples.

LEARN MORE ABOUT
ZIRCLEAN



Core Flo™ DC & Core Flo™ DC Lite

Dual-Cured Core Build-Up Materials

Core-Flo DC and Core Flo DC Lite are dual-cured, core products that are ideal for core build-ups, post cementation and as a dentin replacement material.



High Compressive Strength

Provides high compressive strength in LC and SC modes



High Flexual Strength

Offers high flexual strength in LC and SC modes



Radiopaque

Easy to identify on radiographs for quick and effective diagnosis



Cuts Like Dentin

Allows for easy preparation



Stackable (Core-Flo DC)

Optimal handling when building up core



Self-Leveling (Core-Flo DC Lite)

Allows for excellent adaption resulting in gap-free margins

Dentistry courtesy of Ross Nash, DDS, FAACD



1. Enamel etched with Select HV Etch



2. Core-Flo DC applied after use of Universal Primer™



3. Core-Flo DC placed

LEARN MORE ABOUT
CORE-FLO DC +
CORE-FLO DC LITE



Order Info: Core-Flo DC Lite

System Kit with Universal Primer
Natural/A1 Dual-Syringe (8g)
Opaque White Dual-Syringe (8g)

AB-17410K
A-17801P
A-17803P

Core-Flo DC

Natural/A1 Dual-Syringe (8g)
Opaque White Dual-Syringe (8g)

A-23011P
A-23012P

Accessories

Dispenser
Endo Mixing Tips
Intraoral Mixing Tips

L-22020P
X-81267P
X-81257P





TECHNIQUES

Self-Etch, Total-Etch, and Selective-Etch Bonding	Page 16
Pulp Exposures (Direct Pulp Capping)	Page 17
Deep Preparations (Indirect Pulp Capping)	Page 17
Sandwich Technique	Page 18
Pulpotomy	Page 18
Core Build-ups	Page 19
Intraoral Repairs of Porcelain (PFM or PFZr)	Page 20
Conventional Cementation of Crowns (Retentive Preps)	Page 21
Bonded Cementation of Inlays/Onlays, Bridges, Crowns (Short/Tapered Preps)	Page 22
Veneer Cementation	Page 23

QUICK GUIDES

Zirconia Bonding & Sandwich Technique	Page 24
Veneer Cementation & Cementation With a Bonding Agent	Page 25
Cementation Without a Bonding Agent & Post Cementation With Core Build-up	Page 26



Self-Etch Technique



No phosphoric acid applied to the tooth structure.

Total-Etch Technique



Phosphoric acid (such as Uni-Etch® w/ BAC) applied to both enamel and dentin surface.

Selective-Etch Technique



Phosphoric acid (such as Select HV® w/ BAC) applied to the enamel surface only.

Bonding Technique* using All-Bond Universal



1. Apply two separate coats of All-Bond Universal®, scrubbing the preparation with a microbrush for 10-15 seconds per coat. Do not light cure between coats.



2. Evaporate solvent by thoroughly air-drying with an air syringe for at least 10 seconds; there should be no visible movement of the material. The surface should have a uniform glossy appearance; otherwise, repeat Step 1-2. Light cure for 10 seconds.



3. Continue with placement of the restorative material according to the manufacturer's instructions.

* Refer to instructions for complete details.

Pulp Exposures (Direct Pulp Capping)*

with TheraCal LC



1. Under rubber dam isolation, complete cavity preparation.



2. Achieve hemostasis. Leave dentin visibly moist.



3. Apply TheraCal LC* directly to the exposed pulp. Layer is not to exceed 1 mm in depth. Cover all the exposed areas and extend TheraCal LC at least 1mm onto sound dentin surrounding the exposure.



4. Light cure for 20 seconds. Place desired adhesive, base, and/or restoration following manufacturer's directions. Continue restoring tooth.

Deep Preparations (Indirect Pulp Capping)*

with TheraCal LC



1. Isolate the tooth and perform conventional cavity preparation. Remove all infected carious tooth structure. Leave dentin visibly moist.



2. Apply TheraCal LC directly to the cavity floor of the preparation. Layer is not to exceed 1 mm in depth. Manipulate into a smooth surface covering all deep dentin areas.



3. Light cure for 20 seconds. Place desired adhesive, base, and/or restoration following manufacturer's directions. Continue restoring tooth.




* Refer to instructions for complete details.

Sandwich Technique*

using TheraBase




1. Prepare cavity. Clean the preparation with pumice and water. Rinse thoroughly and dry.



2. Apply TheraBase® to the dentin surfaces of the prepared cavity directly from the syringe.



3. Light cure for 20 seconds or allow to self-cure.



4. Proceed with bonding procedure, such as All-Bond Universal® following manufacturer's instructions.




5. Restore with composite following manufacturer's instructions.

Pulpotomy Treatment*


using TheraCal PT




1. Perform pulpotomy treatment.



2. Achieve hemostasis.



3. Place TheraCal PT® directly in the pulp chamber, and ensure good adaptation to the cavity walls and margins.



4. Light cure TheraCal PT for 10 seconds. Place desired adhesive, base, and/or restoration following manufacturer's directions for restoring tooth.

* Refer to instructions for complete details.

Core Build-Ups*


with Core-Flo DC



1. Isolate tooth and prepare accordingly. Rinse and dry.



2. Dispense 1-2 drops of Universal Primer™ A & B into a clean well. Apply two separate coats of Universal Primer, scrubbing the preparation with a microbrush for 10-15 seconds per coat.



3. Evaporate excess solvent by thoroughly air-drying with an air syringe for at least 10 seconds; there should be no visible movement of the adhesive. The surface should have a uniform glossy appearance; otherwise, apply an additional coat of Universal Primer A & B and air dry.



4. Place the end of the intra-oral tip directly onto the preparation, and express Core-Flo™ DC or Core-Flo™ DC Lite.



5. Light cure the Core-Flo DC or Core-Flo DC Lite for 40 seconds and continue with final preparation.




TO LEARN MORE ABOUT
UNIVERSAL PRIMER A + B
VISIT WWW.BISCO.COM

* Refer to instructions for complete details.

Intraoral Repairs of Porcelain (PFM, or PFZr)*

1. Isolate the area to be repaired. Remove the glaze and bevel (45 degrees) the porcelain around the area to be repaired. Sandblast or abrade with a coarse diamond bur. Rinse with water and air dry.



2. Place Barrier Gel on the gingival tissue and porcelain to protect areas where etching is not desired.



3. Apply 9.5% Porcelain Etchant to the dry porcelain surface for 90 seconds. Suction the etchant with high volume evacuation, then rinse with water and air dry. The etched porcelain should appear dull and frosty.



4. Apply 1 coat of silane (Porcelain Primer) to the etched porcelain and allow to dwell for 30 seconds. Dry with (warm) air syringe.



5. Apply 1 coat of Z-Prime™ Plus to the exposed metal/zirconia/alumina and dry with an air syringe for 3-5 seconds.



6. If metal masking is required, use Dual-Cured Opaquer Base and Catalyst, one drop of each, onto a mixing pad and mix with a brush tip. Apply a thin coat of the mixture only to the metal surface. Allow to self cure, or light cure for 5 seconds.



7. Apply a thin layer of Porcelain Bonding Resin to the repair site. Apply a hybrid composite (Aelite™ All-Purpose Body) to replace fractured porcelain and light cure, finish/polish.

LEARN MORE ABOUT
THE INTRAORAL
REPAIR KIT




* Refer to instructions for complete details.


Conventional Cementation of Crowns (Retentive Preps)* using TheraCem



1. Sandblast internal surface of restoration (unless restoration has previously been sandblasted).



2. Remove temporary restoration and clean prep with pumice and water slurry. Rinse thoroughly and dry.



3. Try-in (to check occlusion and fit). After try-in, thoroughly rinse the restoration with water spray and dry.



4. Outside of the mouth, cover all bonded surfaces of the restoration with a layer of ZirClean®. Allow 20 seconds for the cleaning action of ZirClean to take affect, then thoroughly rinse with water spray and dry.



5. Dispense TheraCem® into the restoration. Seat the restoration with gentle passive pressure and remove any excess cement.

PRO TIP: To aid in the removal of excess cement, initially light-cure the margins for 2-3 seconds.



6. Light cure for 20-30 seconds or allow to self cure.

* Refer to instructions for complete details.

Bonded Cementation of Inlay/Onlays, Bridges, Crowns (Short/Tapered Preps)* using Duo-Link Universal




1a. Porcelain/Lithium Disilicate: Apply a thin coat of silane (Porcelain Primer) to the internal surface. Wait for 30 seconds, or dry with (warm) air.


OR



1b. If restoration is metal/zirconia/indirect composite sandblast internal surface of restoration, unless restoration has previously been sandblasted.



2. Remove temporary restoration and clean prep with pumice and water slurry. Rinse thoroughly and dry.



3. Try-in (to check occlusion and fit). After try-in, thoroughly rinse the restoration with water spray and dry.



4. Outside of the mouth, Cover all bonded surfaces of the restoration with a layer of ZirClean®. Allow 20 seconds for the cleaning action of ZirClean to take affect, then thoroughly rinse with water spray and dry.



5. If restoration is metal/zirconia/indirect composite, apply one coat of Z-Prime™ Plus to the internal surface of the restoration and air dry for 3-5 seconds.



6. If prep includes enamel, selectively etch enamel with Select HV® Etch w/BAC for 15 seconds, suction then rinse thoroughly.



7. Dispense Universal Primer™ and mix (or All-Bond Universal®) in a mixing well. Apply 2 separate coats, agitating each coat for 10-15 seconds.



8. Gently air dry until there is no visible movement of the adhesive. Then thoroughly air dry with greater air pressure. The surface should appear shiny, otherwise repeat step 7. Light cure for 10 seconds.**



9. Using a dual-cured resin cement (Duo-Link Universal™), fill the internal surface of the restoration and/or the prep with the cement. Seat with gentle, passive pressure and remove excess cement while holding restoration in place. Allow the cement to self-cure, or light-cure each surface of the tooth for 40 seconds.

* Refer to instructions for complete details.

** Universal Primer does not require light cure.



Veneer Cementation* using Choice™ 2



OR



1. Prepare the internal surface of the restoration as instructed by the laboratory:

a. Porcelain/Lithium Disilicate:
If needed, etch the veneer with hydrofluoric acid (4% Porcelain Etchant or 9.5% Porcelain Etchant) according to the manufacturer's instructions.

b. Apply a thin coat of silane (Porcelain Primer/Bis-Silane™) to the internal surface. Wait for 30 seconds, or dry with (warm) air.



2. Remove the temporary restorations and clean the preparations (Cavity Cleanser™ and pumice slurry). Rinse thoroughly.



3. Try in the restorations using the corresponding shade of water-soluble try-in paste (Choice™ 2 Try-In Paste). Remove the veneers and either clean with etchant (Uni-Etch® w/BAC or Select HV® Etch w/BAC) and rinse thoroughly, or ultrasonicate in water or alcohol for 2-3 minutes. Dry the restorations.



4. Isolate. Etch preparations (Uni-Etch w/BAC or Select HV Etch w/BAC) for 15 seconds, rinse thoroughly.



5. Dispense adhesive (All-Bond Universal®) in a mixing well. Apply 2 separate coats, agitating each coat for 10-15 seconds.



6. Gently air dry until there is no visible movement of the adhesive. Then thoroughly air dry with greater air pressure. The surface should appear shiny, otherwise apply additional coats.



7. Light cure for 10 seconds.



8. Apply a thin layer of HEMA-free resin (Porcelain Bonding Resin) to the internal surface of the veneer. Do NOT light cure.



9. Using a light-cured resin cement (Choice 2), fill the internal surface of the restoration with the cement.




10. Seat with gentle, passive pressure and light cure for 2-3 seconds to tack the veneer into place. Remove excess cement, then light cure each veneer for 40 seconds.

* Refer to instructions for complete details.

Zirconia Bonding




Sandwich Technique




Veneer Cementation




Cementation With a Bonding Agent



Cementation Without a Bonding Agent



Post Cementation With Core Build-up



BISCO's Direct-to-You Promise

BISCO manufactures, sells & ships products **Direct-To-You**, providing you with advantages that you can't get anywhere else. This is our **Direct-to-You** promise...

1. You always receive the **freshest, practically made-to-order products** directly from us, providing you with the longest shelf life on your purchase. We feel you deserve to get the most out of the BISCO products you use.
2. Our **experts are on standby to help you with all your restorative questions.** No automated phone systems - just REAL people, providing REAL help, in REAL time. Looking for advice, tips, or just someone to talk to? We've got you covered!
3. You always receive **Direct-to-You** pricing. We provide top **quality, award-winning, restorative products at competitive prices,** without a distributor markup. In other words, you can have your cake and eat it too... REALLY!
4. **We stand behind the quality and integrity of our products.** If, for any reason, you are not satisfied with your purchase, we will give you a 100% refund.
5. At BISCO, you are not just a customer, **you are family.** We love connecting with our customers and hearing what you have to say. We believe it is this connection with you that allows us to discover new and innovative ways to solve the restorative problems you face every day in your office.
6. BISCO is a **family-owned company** located in Schaumburg, Illinois. All our products are made by us, here in the U.S.A.
7. **We are here for you,** and we want to help you perform your BEST dentistry. When you succeed, we succeed, and ultimately, your patients benefit... everyone wins!

All this and more . . . DIRECT TO YOU!



BISCO, Inc.
1100 W. Irving Park Rd.
Schaumburg, IL 60193, USA
1-800-247-3368
www.bisco.com

Visit us at www.bisco.com
for a complete product catalog!
Rx only.

